

The two-hour orbit of a binary millisecond X-ray pulsar

Deepto Chakrabarty & Edward H. Morgan

Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139 USA

Typical radio pulsars are magnetized neutron stars that are born rapidly rotating and slow down as they age on time scales of 10 to 100 million years. However, millisecond radio pulsars spin very rapidly even though many are billions of years old.¹ The most compelling explanation is that they have been “spun up” by the transfer of angular momentum during accretion of material from a companion star in so-called low-mass X-ray binary systems, LMXBs. (LMXBs consist of a neutron star or black hole accreting from a companion less than one solar mass.²) The recent detection of coherent X-ray pulsations with a millisecond period from a suspected low-mass X-ray binary system appears to confirm this link.³ Here we report observations showing that the orbital period of this binary system is two hours, which establishes it as an LMXB. We also find an apparent modulation of the X-ray flux at the orbital period (at the two per cent level), with a broad minimum when the pulsar is behind this low-mass companion star. This system seems closely related to the “black widow” millisecond radio pulsars, which are evaporating their companions through irradiation.^{4–8} It may appear as an eclipsing radio pulsar during periods of X-ray quiescence.

The transient X-ray source SAX J1808.4–3658 was first observed in September 1996 by the Wide Field Cameras on the BeppoSAX X-ray satellite during a bright state which lasted about 20 days.⁹ The source was not detected (X-ray flux $< 3 \times 10^{-11}$ erg cm⁻² s⁻¹ in the 2–10 keV band) during an August 1996 observation, but reached a peak X-ray flux of 2×10^{-9} erg cm⁻² s⁻¹ during the September bright state. Also during the bright state, BeppoSAX detected two type I X-ray bursts from the source, each lasting less than 30 s. Such bursts are due to thermonuclear ignition of accreted material on the surface of neutron stars that have low ($< 10^{10}$ G) magnetic fields.¹⁰ Analysis of the bursts in SAX J1808.4–3658 indicates that it is 4 kpc distant and has a peak X-ray luminosity of 6×10^{36} erg s⁻¹ in its bright state and $< 10^{35}$ erg s⁻¹ in quiescence.⁹ The source position was determined to within a 1σ radius of 0.6 arcmin.

More recently, a serendipitous slew of the Proportional Counter Array (PCA) on the Rossi X-Ray Timing Explorer (RXTE) over this region of the sky on 9 April 1998 indicated the presence of an X-ray source, designated XTE J1808–369. Repeated scans over the region with the PCA on 11 April localized the position with sufficient precision to confirm that the source was probably the same as SAX J1808.4–3658.¹¹ RXTE subsequently made numerous pointed observations of the source during 11 April–6 May on a public “target-of-opportunity” basis. The source reached a peak luminosity of 5×10^{36} erg s⁻¹ on 13 April and had faded below 10^{35} erg s⁻¹ by 6 May.

Strong coherent 400 Hz pulsations were clearly detected in the 2–30 keV PCA data from most of these observations, and were first reported by Wijnands and van der Klis.^{12,3} These are the fastest persistent coherent pulsations ever detected from an X-ray binary. From standard magnetic disk accretion theory, the detection of X-ray pulsations at a luminosity $\sim 10^{36}$ erg s⁻¹ requires that the neutron star’s surface dipole magnetic field be $B < 10^8$ G (ref. 3). This is much weaker than the $\sim 10^{12}$ G fields found in other (slower) accretion-powered pulsars,¹³ but it is consistent with the weak fields expected for type I X-ray bursts.¹⁰ If pulsations are detected when the source is at lower luminosities, the implied surface field would be even weaker.

In order to search for orbital Doppler shifts, we first selected the 3–30 keV photon arrival times during 11–18 April and binned them into 2^{-11} s (≈ 0.5 ms) samples, after correcting these times for RXTE’s motion with respect to the solar system barycenter using the best BeppoSAX position for the source⁹ (right ascension 18h 08m 29s, declination $-36^\circ 58.6'$, equinox J2000.0). We then measured the barycentric pulse frequency at 128 s intervals using an oversampled Fourier power spectrum. The sequence of pulse frequencies showed an obvious 2 hr sinusoidal modulation which was well fit by a constant spin frequency plus a circular, Keplerian orbit.¹⁴

Using this provisional timing model, we epoch-folded each 128 s interval in the 11–18 April data and cross-correlated the resulting pulse profiles with a sinusoid to measure the pulse phase history, which generally yields more precise timing parameters than a pulse frequency

Table 1. Observed parameters of SAX J1808.4–3658

Barycentric pulse frequency, ν_0 (Hz)	400.9752106(8)
Pulse frequency derivative, $ \dot{\nu} $ (Hz s $^{-1}$)	$< 7 \times 10^{-13}$
Projected semimajor axis, $a_1 \sin i$ (lt-ms)	62.809(1)
Orbital period, P_{orb} (s)	7249.119(1)
Epoch of 90° mean longitude, $T_{\pi/2}$ (MJD)	50914.899440(1)
Eccentricity, e	$< 5 \times 10^{-4}$
Pulsar mass function, f_1 ($M_\odot$)	$3.7789(2) \times 10^{-5}$

Modified Julian date MJD = JD – 2400000.5. Numbers in parentheses are the 1σ uncertainties in the last significant figure, and upper limits are quoted at the 2σ level. The $\dot{\nu}$ limit refers to the magnitude, independent of sign. Epochs are given in Barycentric Dynamical Time (TDB).

history. These pulse phases were well fit (reduced $\chi^2=1.26$ with 441 degrees of freedom) by a constant spin frequency ν plus a circular Keplerian orbit. The best-fit parameters are given in Table 1. The pulse arrival time delays with respect to a constant frequency model are plotted in Figure 1. No improvement in the fit to the 11–18 April data is obtained by including an orbital eccentricity e or a spin frequency derivative $\dot{\nu}$. A more sensitive search for a non-zero value of $\dot{\nu}$, which includes the later data taken as the X-ray outburst faded from view, will be presented elsewhere.

The present limits on $\dot{\nu}$ constrain the positional uncertainty of the star. An error of 0.6 arcmin in the assumed source position would give rise to periodic timing residuals with 1 yr period and ~ 80 ms amplitude,¹⁵ and such residuals would lead to an apparent $\dot{\nu}$ in the timing solution for observations shorter than 3 months. Our limit on $\dot{\nu}$ from the 11–18 April data suggests that the BeppoSAX position is accurate to within 25 arcsec in ecliptic longitude. We note that a probable optical counterpart has been identified, lying 19 arcsec from the BeppoSAX position.^{16,17}

The pulsar mass function f_1 , which relates the pulsar mass m_1 , the companion mass m_2 , and the binary inclination i (the angle between the line of sight and the orbital angular momentum vector), may be computed from the observed Keplerian parameters $a_1 \sin i$ and P_{orb} ,

$$f_1 \equiv \frac{(m_2 \sin i)^3}{(m_1 + m_2)^2} = \frac{4\pi^2 (a_1 \sin i)^3}{GP_{\text{orb}}^2}, \quad (1)$$

from which we find $f_1 = 3.7789(2) \times 10^{-5} M_\odot$ (here $M_\odot$ is the solar mass). Given assumed values of m_1 and i , equation (1) can be solved for m_2 . For m_1 , dynamical mass measurements in both binary millisecond radio pulsars¹⁸ and accretion-powered pulsars¹⁹ are all generally consistent with a neutron star mass of $1.35 M_\odot$. However, neutron stars in LMXBs have undergone substantial accretion from their binary companion and may be as massive as $2 M_\odot$ (e.g., ref. 20). For i , we note that for an ensemble of binaries whose inclinations are distributed randomly, the *a priori* probability of observing a system with inclination i or smaller is $(1 - \cos i)$. If SAX J1808.4–3658 is drawn




Figure 1. Pulse arrival time delays due to the 2 hr binary orbit, with respect to a constant pulse frequency model. Mean longitude is the orbital phase angle measured from the ascending node, so that the pulsar is behind the companion at 90°. The squares are the measured time delays, the solid curve is the best-fit orbital model, and the diamonds are the model residuals (on a 50× expanded scale). The residuals have an r.m.s. uncertainty of 75 μ s, shown for one of the points.

from such an ensemble, then there is a 95% probability that $m_2 < 0.14 M_\odot$ if $m_1 = 1.35 M_\odot$. For $m_1 = 2 M_\odot$, we find $m_2 < 0.18 M_\odot$. In either case, the binary separation is most likely of order 1 lt-s.

The nature of the companion star provides an important clue to the evolutionary history of an LMXB. The only direct information we have on the companion comes from observations of the probable optical counterpart, which suggest that the companion is a faint low-mass star subject to X-ray heating by the pulsar.^{16,17} We can use the binary parameters to deduce more about this star. The companions in most LMXBs fill (or nearly fill) their critical gravitational potential surface, known as the Roche lobe. Because the mean density of a Roche-lobe-filling companion (for the case where $m_2 < m_1$) is uniquely determined by the binary period,²¹ we can relate the companion’s radius in SAX J1808.8–3658 to its mass as $R_2 = 0.17 R_\odot (m_2/0.1 M_\odot)^{1/3}$, giving a minimum companion radius of $0.12 R_\odot$ (here $R_\odot$ is the solar radius). The assumption of Roche lobe overflow thus strongly constrains the nature of the companion. Given the compact size of the binary, only white dwarf (WD) or low-mass main sequence (MS) companions are plausible. (In principle, a helium-burning star could also fit in this binary, but we would expect it to be much brighter than the proposed optical counterpart.²²) A WD with mass m has radius $R = 0.013 R_\odot (1 + X)^{5/3} (m/M_\odot)^{-1/3}$, where X is the hydrogen mass fraction.²³ Comparing this to our equation for R_2 , we see that a helium WD ($X = 0$) would be too

small, and hence is ruled out. Even a hydrogen-rich WD ($X = 0.9$) is essentially excluded for all but $i \approx 90^\circ$. However, the lack of a deep X-ray eclipse rules out $i > 82^\circ$, assuming a Roche-lobe-filling companion.

Normal low-mass hydrogen MS stars have $R/R_\odot \approx m/M_\odot$, and comparison of detailed models²⁴ with our equation for R_2 yields a Roche-lobe-filling mass of $0.17 M_\odot$. This would require a small binary inclination ($i < 20^\circ$), which has low *a priori* probability (5%). However, normal MS models are inappropriate in our case, since the stellar structure must be strongly influenced by irradiation from the pulsar. If the incident flux at the companion surface exceeds $\sim 10^{10} \text{ erg cm}^{-2} \text{ s}^{-1}$, the star may be “bloated” to larger radii.²⁵ This is especially true for MS stars having $m < 0.3 M_\odot$, which have fully convective envelopes. For the ~ 1 lt-s binary separation in this system, the X-ray flux at the companion will exceed $10^{14} \text{ erg cm}^{-2} \text{ s}^{-1}$, and the bloating can be severe. Under these conditions, MS stars as light as $0.1 M_\odot$ or less may fill their Roche lobe, consistent with a more likely inclination.

Another possibility is that irradiation by the neutron star might directly drive mass loss from the companion (“ablation”) whether or not it fills its Roche lobe.²⁶ The binary parameters of SAX J1808.4–3658 are very similar to those of the five “black widow” millisecond radio pulsars in very close binaries, all of which are ablating their low mass companions.^{4–8} At least four of these radio systems show eclipses which are too broad to be caused by a Roche-lobe-filling companion, and which are instead attributed to an ablated wind.^{4–6,8}

We find evidence that the X-ray flux from SAX J1808.4–3658 is slightly modulated at the orbital period. The apparent modulation is roughly sinusoidal with 2% amplitude and a minimum when the pulsar is behind the companion, and it is much broader and much shallower than the (at most) 5 minute eclipse possible from a Roche-lobe-filling companion. However, the detailed features of this 2 hr modulation must be viewed with caution, since its strength is comparable to the variation of the RXTE background with the 96 min spacecraft orbit. The similarity in strength and time scales of these two variations makes them difficult to disentangle, and final confirmation must await either additional data or an improvement in the RXTE background model. Still, the detection of a 2 hr source flux modulation (if not its precise morphology) seems secure.

We suggest that the intensity dips are due to scattering in an ablated wind. Given the similarity between this source and the eclipsing radio pulsars, SAX J1808.4–3658 may emerge as a radio pulsar during X-ray quiescence.²⁷ Moreover, since the radio emission would be less penetrating than the X-rays, a deeper eclipse dip might be observed, possibly providing a strong constraint on the binary inclination. The presence of the slight X-ray dips, if confirmed, would rule out small binary inclinations and

make it highly likely that the companion mass is less than $0.1 M_\odot$.

Whether the mass accretion is fed by Roche-lobe overflow or an ablated wind or both, we can understand the transient nature of the X-ray emission as long as some of this material forms an accretion disk. Although the instantaneous mass accretion rate during the 1996 and 1998 bright states was $\approx 3 \times 10^{-10} M_\odot \text{ yr}^{-1}$, the small duty cycle of the X-ray activity indicates a long-term mean mass transfer rate of $\dot{M} \approx 1 \times 10^{-11} M_\odot \text{ yr}^{-1}$. For such a low $\dot{M}$, the accretion disk would probably be subject to dwarf-nova-type instabilities, leading to episodic outbursts of X-ray emission.^{28,29} We note that for a Roche-lobe-filling companion, angular momentum losses due to gravitational radiation would drive a mass transfer rate of $\approx 10^{-11} M_\odot \text{ yr}^{-1}$ for a $0.05 M_\odot$ secondary,³⁰ consistent with our inferred inclination constraints.

Future X-ray outbursts from SAX J1808.4–3658 may allow detection of a spin frequency derivative, which would further constrain both the mass accretion rate and the pulsar’s magnetic field strength and would provide a probe of magnetic disk accretion torque theory in the previously unexplored regime of a very small magnetosphere. The characteristic accretion torque expected for the observed X-ray luminosity should cause a $\dot{\nu}$ of order $10^{-14} \text{ Hz s}^{-1}$, which is detectable in observations spanning a month or more. Further observations may also detect orbital period evolution, which would probe the competing effects of gravitational radiation, tidal interactions, and mass loss on the binary angular momentum. An astrophysically interesting limit of $|\dot{P}_{\text{orb}}| < 10^{-11}$ could be reached in less than a year of observations.

Received 23 April 1998; Accepted 16 June 1998.

1. Backer, D. C., Kulkarni, S. R., Heiles, C., Davis, M. M. & Goss, W. M. A millisecond pulsar, *Nature* **300**, 615–618 (1982).
2. Bhattacharya, D. & van den Heuvel, E. P. J. Formation and evolution of binary and millisecond pulsars, *Phys. Reports* **203**, 1–124 (1991).
3. Wijnands, R. & van der Klis, M. Discovery of the first accretion-powered millisecond X-ray pulsar, *Nature*, submitted (astro-ph/9804216) (1998).
4. Fruchter, A. S. *et al.* The eclipsing millisecond pulsar PSR 1957+20, *Astrophys. J.* **351**, 642–650 (1990).
5. Nice, D. J. & Thorsett, S. E. Pulsar PSR 1744–24A: timing, eclipses, and the evolution of neutron star binaries, *Astrophys. J.* **397**, 249–259 (1992).
6. Manchester, R. N. *et al.* Discovery of ten millisecond pulsars in the globular cluster 47 Tucanae, *Nature* **352**, 219–221 (1991).
7. Deich, W. T. S. *et al.* The binary pulsar PSR 1908+00 in NGC 6760, *Astrophys. J.* **410**, L95–L98 (1993).
8. Stappers, B. W. *et al.* Probing the eclipse region of a binary millisecond pulsar, *Astrophys. J.* **465**, L119–L122 (1996).

9. in 't Zand, J. J. *et al.* Discovery of the X-ray transient SAX J1808.4–3658, a likely low-mass X-ray binary, *Astr. Astrophys.* **331**, L25–L28 (1998).
10. Lewin, W. H. G., van Paradijs, J. & Taam, R. E. in *X-Ray Binaries* (eds Lewin, W. H. G., van Paradijs, J. & van den Heuvel, E. P. J.) 175–232 (Cambridge Univ. Press, Cambridge, 1995).
11. Marshall, F. E. SAX J1808.4–3658 = XTE J1808–369, *IAU Circ.*, No. 6876 (1998).
12. Wijnands, R. & van der Klis, M. SAX J1808.4–3658 = XTE J1808–369, *IAU Circ.*, No. 6876 (1998).
13. White, N. E., Nagase, F. & Parmar, A. N. in *X-Ray Binaries* (eds Lewin, W. H. G., van Paradijs, J. & van den Heuvel, E. P. J.) 1–57 (Cambridge Univ. Press, Cambridge, 1995).
14. Chakrabarty, D. & Morgan, E. H. SAX J1808.4–3658 = XTE J1808–369, *IAU Circ.*, No. 6877 (1998).
15. Lyne, A. G. & Graham-Smith, F. *Pulsar Astronomy*. (Cambridge U. Press, Cambridge, 1990).
16. Roche, P. *et al.* SAX J1808.4–3658 = XTE J1808–369, *IAU Circ.*, No. 6885 (1998).
17. Giles, A. B., Hill, K. M. & Greenhill, J. G. SAX J1808.4–3658 = XTE J1808–369, *IAU Circ.*, No. 6886 (1998).
18. Thorsett, S. E. & Chakrabarty, D. Neutron star mass measurements. I. Radio pulsars., *Astrophys. J.*, submitted (astro-ph/9803260) (1998).
19. van Kerkwijk, M. H., van Paradijs, J. & Zuiderwijk, E. J. On the masses of neutron stars, *Astr. Astrophys.* **303**, 497–501 (1995).
20. Zhang, W., Strohmayer, T. E. & Swank, J. H. Neutron star masses and radii as inferred from kilohertz quasi-periodic oscillations, *Astrophys. J.*, L167–L170 (1997).
21. Faulkner, J., Flannery, B. P. & Warner, B. Ultrashort-period binaries. II. HZ 29 (=AM CVn): a double-white-dwarf semidetached postcataclysmic nova?, *Astrophys. J.* **175**, L79–L83 (1972).
22. Savonije, G. J., de Kool, M. & van den Heuvel, E. P. J. The minimum orbital period for ultra-compact binaries with helium burning secondaries, *Astr. Astrophys.* **155**, 51–57 (1986).
23. Paczynski, B. Gravitational waves and the evolution of close binaries, *Acta Astron.* **17**, 287–296 (1967).
24. Tout, C. A., Pols, O. R., Eggleton, P. P. & Han, Z. Zero-age main-sequence radii and luminosities as analytic functions of mass and metallicity, *Mon. Not. R. astr. Soc.* **281**, 257–262 (1996).
25. Podsiadlowski, P. Irradiation-driven mass transfer in low-mass X-ray binaries, *Nature* **350**, 136–138 (1991).
26. D’Antona, F. in *Evolutionary Processes in Binary Stars* (eds Wijers, R. A. M. J., Davies, M. B. & Tout, C. A.) 287–306 (Kluwer, Dordrecht, 1996).
27. Stella, L., Campana, S., Colpi, M., Mereghetti, S. & Tavani, M. Do quiescent soft X-ray transients contain millisecond radio pulsars?, *Astrophys. J.* **423**, L47–L50 (1994).
28. van Paradijs, J. On the accretion instability in soft X-ray transients, *Astrophys. J.* **464**, L139–L141 (1996).
29. King, A. R., Kolb, U. & Burderi, L. Black hole binaries and X-ray transients, *Astrophys. J.* **464**, L127–L130 (1996).
30. Verbunt, F. & van den Heuvel, E. P. J. in *X-Ray Binaries* (eds Lewin, W. H. G., van Paradijs, J. & van den Heuvel, E. P. J.) 457–494 (Cambridge Univ. Press, Cambridge, 1995).

Acknowledgements. We thank L. Bildsten, V. Kaspi, A. Levine, R. Nelson, R. Remillard, F. Rasio, S. Thorsett, M. van der Klis, and B. Vaughan for useful discussions, and M. Muno for assistance with the data analysis. We also thank F. Marshall, J. Swank and the RXTE team at NASA/Goddard Space Flight Center for arranging these target-of-opportunity observations and the necessary follow-up. This work was supported by NASA.

Correspondence should be addressed to D.C. (e-mail: deepto@space.mit.edu).